

GSP . Jan 27

On certain Lagrangian subvarieties in minimal resolution of Kleinian singularities

- 1) Reminder on Kleinian singularities
- 2) Anti-Poisson involution & their fixed loci
- 3) Preimage of fixed loci under minimal resolution

Overview:

minimal resolution

$$\begin{array}{c} \tilde{X} \\ \cup \\ \pi^{-1}(X^\theta) \end{array}$$

$$\xrightarrow{\pi}$$

Kleinian singularity

$$\begin{array}{c} X \\ \cup \\ X^\theta \end{array}$$

anti-Poisson involution

(singular) Lagrangian subvariety.

Goal: Describe X^θ and $\pi^{-1}(X^\theta)$ as schemes. $\left\{ \begin{array}{l} \text{irreducible component} \\ \text{intersection pattern} \\ \text{reduced or not (multiplicity)} \end{array} \right.$

Motivation: Classification of irreducible $HC(\mathfrak{g}, K)$ -mod

• G simple algebraic group; \mathfrak{g} \mathcal{N} .

$\sigma: \mathfrak{g} \rightarrow \mathfrak{g}$, Lie algebra involution $\leadsto \mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}$.

K , corresponding connected alg group

A $HC(\mathfrak{g}, K)$ -module is a f.g. $U(\mathfrak{g})$ -mod, M , s.t. $K \curvearrowright M$

locally finitely & integrates to $K \curvearrowright M$.

associated variety $AV(M) = \text{Supp}(gr M)$

M irreducible $\leadsto AV(M)$ is finite union of K -orbits in $\mathcal{N} \cup \mathfrak{p}$

• Take $\mathcal{O} \subset \mathcal{N}$ nilpotent orbit, J unipotent ideal associated to \mathcal{O}

• $\text{codim}_{\mathbb{C}}(\partial \bar{\mathcal{O}} \cdot \bar{\mathcal{O}}) \geq 4$

\mathcal{O}_K is a K -orbit in $\mathcal{O} \cap \mathfrak{p}$.

[Losev & Tu, 23] classified irred M s.t. $J \subset \text{Ann}(M)$ & $AV(M) = \overline{\mathcal{O}_K}$

• $\text{codim}_{\mathbb{C}}(\partial \bar{\mathcal{O}} \cdot \bar{\mathcal{O}}) = 2$, classification unknown!

Take $\mathcal{O}' \subset \bar{\mathcal{O}}$ s.t. $\text{codim}_{\mathbb{C}}(\partial \bar{\mathcal{O}} \cdot \bar{\mathcal{O}}) = 2$

Take $e' \in \mathcal{O}'$, e' a normal point in $\bar{\mathcal{O}} \rightarrow S^1$, Slodowy slice $\leadsto S^1 \cdot \bar{\mathcal{O}} \simeq X$, a Kleinian singularity

$\mathcal{O}' \curvearrowright S^1 \cdot \bar{\mathcal{O}} \leadsto \underbrace{S^1 \cdot \bar{\mathcal{O}} \cap \mathfrak{p}} = (S^1 \cdot \bar{\mathcal{O}})^{\mathcal{O}} \simeq X^{\mathcal{O}}$, fixed loci

$\mathcal{O} := -\sigma$, $\mathfrak{p} := \mathfrak{g}^{\mathcal{O}}$

anti-Poisson involution.

local picture of $AV(M)$ near e'

§ 1. Reminder on Kleinian singularities

Let $\Gamma \subset \text{SL}_2(\mathbb{C})$ be a finite subgroup.

Kleinian singularity $X := \mathbb{C}^2/\Gamma = \text{Spec } \mathbb{C}[u, v]^\Gamma$

Example: $\Gamma = \{\pm I_2\}$ (A_1)


$$\mathbb{C}[u, v]^\Gamma = \text{even degree polynomials} = \mathbb{C}[x=u^2, y=v^2, z=uv] \\ = \mathbb{C}[x, y, z] / (xy - z^2)$$

Theorem (Klein, 1884): $X = \mathbb{C}^2/\Gamma \hookrightarrow \mathbb{C}^3$ (single relation) with an isolated singularity at 0.

$\pi: \tilde{X} \rightarrow X$, minimal resolution

exceptional locus $\pi^{-1}(0)_{\text{red}} = C_1 \cup \dots \cup C_n$ ($C_i \cong \mathbb{P}^1$)
irreducible components

Dual graph of $\pi^{-1}(0)_{\text{red}}$:



Fact (Du Val, 1934): Dual graphs of $\pi^{-1}(0)_{\text{red}}$ are ADE Dynkin diagrams. (bijection)

Rem: $\pi^{-1}(0)$ is not reduced in general!


\mathfrak{g} : simple Lie algebra of types ADE, simple root system $\{\alpha_1, \alpha_2, \dots, \alpha_n\}$
 δ - unique maximal root, $\delta = \sum_{i=1}^n m_i \alpha_i$ in the adjoint rep $\mathfrak{g} \cong \mathfrak{g}$.

Then we have $\pi^{-1}(0) = \sum_{i=1}^n m_i C_i$ as a divisor. [Artin 66].

Type A_n : $\alpha_i = \epsilon_i - \epsilon_{i+1}$, $\delta = \epsilon_1 - \epsilon_{n+1} = \alpha_1 + \dots + \alpha_n \rightsquigarrow \pi^{-1}(0)_{\text{reduced}}$


 $(m_1, \dots, m_n) = (1, \dots, 1)$

Type D_n $(m_1, \dots, m_n) = (1, 2, \dots, 2, 1, 1) \rightsquigarrow \pi^{-1}(0)$ not reduced



Examples:


1) A_1



$\pi^{-1}(0)_{red} = \mathbb{P}^1$ dual graph \bullet A_1 Dynkin diagrams

More generally, type A: $\pi^{-1}(0)$ (reduced)

2) D_4



dual graph D_4 Dynkin diagram

§2. Anti-Poisson involution

$\mathbb{C}[u, v]$, Poisson bracket $\{f_1, f_2\} = \frac{\partial f_1}{\partial u} \frac{\partial f_2}{\partial v} - \frac{\partial f_1}{\partial v} \frac{\partial f_2}{\partial u}$ (deg = -2)

$\mathcal{P} \subset SL_2(\mathbb{C}) \rightsquigarrow \mathcal{P}$ preserves $\{, \}$.

$\rightarrow \mathbb{C}[u, v]^{\mathcal{P}} \subset \mathbb{C}[u, v]$, graded Poisson subalgebra (deg = -2)

Example: Type A_n : $\mathbb{C}[X] = \mathbb{C}[x, y, z] / (xy - z^{n+1})$

$$\{x, y\} = (n+1)z^n$$

$$\{x, z\} = (n+1)x$$

$$\{y, z\} = -(n+1)y$$

Def: An anti-Poisson involution of a Kleinian singularity $X \simeq \mathbb{C}^2/p$ is a graded algebra involution $\theta: \mathbb{C}[X] \rightarrow \mathbb{C}[X]$ such that

$$\theta(\{f_1, f_2\}) = -\{\theta(f_1), \theta(f_2)\} \quad \forall f_1, f_2 \in \mathbb{C}[X].$$

Example: Type A_n : $\theta: \mathbb{C}[X] \rightarrow \mathbb{C}[X]$

$$x \mapsto y, \quad y \mapsto x, \quad z \mapsto z$$

is an anti-Poisson involution.

Def (fixed locus):

$$X^\theta := \text{Spec } \mathbb{C}[X] / \mathcal{I}, \quad \text{where } \mathcal{I} = (\theta(f) - f, f \in \mathbb{C}[X])$$

Example (continued)

$$X^\theta = \text{Spec } \mathbb{C}[x, y, z] / (xy - z^{n+1}, x - y) \simeq \text{Spec } \mathbb{C}[x, z] / (x^2 - z^{n+1}) \text{ reduced.}$$

- union of two \mathbb{A}^1 when n odd
- a cusp when n even.

Prop 1 (Classification of θ)

There are finitely many anti-Poisson involutions on \mathbb{C}^2/\mathcal{T} up to conjugation by Poisson automorphisms.

Prop 2 (Description of X^θ)

- (1) X_{red}^θ is either a (singular) Lagrangian subvariety of X , or only contains the singular point.
- (2) X^θ is reduced and connected. Each irreducible component is either an \mathbb{A}^1 or a cusp.

Proof (1) follows from

Exercise

Lemma (M. W) symplectic manifold with anti-symplectic involution τ ($\tau^*\omega = -\omega$)

\rightarrow pf: look at tangent space




Then M^τ is either empty or a Lagrangian submanifold.

applied to $X^{\text{reg}} = X \setminus \{0\}$

(2) follows from classification in Prop 1 + case by case calculations.

Example

Anti-Poisson involutions for Type A_n Kleinian singularity

	case I	case <u>I</u>	case <u>II</u>
<p>A_n, n odd</p> <p>$xy - z^{n+1} = 0$</p> <p>X^0</p> <p>$\pi^{-1}(X^0)$</p>	<p>$x \leftrightarrow y$ </p> 	<p>$z \mapsto -z$ </p> 	<p>$x \mapsto -x$</p> <p>$y \mapsto -y$ </p> <p>$z \mapsto -z$</p> 
<p>A_n, n even</p> <p>$xy - z^{n+1} = 0$</p> <p>X^0</p> <p>$\pi^{-1}(X^0)$</p>	<p>$x \leftrightarrow y$ </p>  <p>the only triple intersection</p>	<p>$x \mapsto -x$</p> <p>$z \mapsto -z$ </p> 	<p>Fixed loci in purple!</p>

Corresponding Cartan involution

outer involution $\sigma: X \mapsto -X^T$

inner involution $\sigma = \text{Ad}(I_{k,e})$

not seen.

D_n, E_6 , two cases of API ; E_7, E_8 one case.

§3 Preimage of fixed loci under minimal resolution

$$\pi: \tilde{X} \rightarrow X, \quad 0 \in X^\theta \rightsquigarrow \pi^{-1}(0) \subset \pi^{-1}(X^\theta)$$

§ 3.1 lift. $\quad \quad \quad : X \rightarrow X$

Def: A lift of $\theta: X \rightarrow X$ is an anti-symplectic involution $\tilde{\theta}: \tilde{X} \rightarrow \tilde{X}$
 s.t. $\pi \circ \tilde{\theta} = \theta \circ \pi$

Thm 3: There exists a unique lift for any anti-Poisson involution θ on X .

Now let's see why the lift helps to determine the preimage, $\pi^{-1}(X^\theta)$

union of \mathbb{P}^1 's

Claim: $\pi^{-1}(X^\theta)_{\text{red}} = \pi^{-1}(0)_{\text{red}} \cup \boxed{\tilde{X}^{\tilde{\theta}}}$ $\tilde{X}^{\tilde{\theta}}$ is smooth Lagrangian (by Lemma)

no intersection! X becomes $||$ in $\pi^{-1}(X^\theta)$;

no cusp! $\{$ becomes $|$ in $\pi^{-1}(X^\theta)$

Let $L \subset X^\theta$ be an irreducible component

define $\tilde{L} := \overline{\pi^{-1}(L \setminus 0)}$, then \tilde{L} is an irreducible component in $\pi^{-1}(X^\theta)$ &

$$\tilde{L} \subset \tilde{X}^{\tilde{\theta}} \Rightarrow \tilde{L} \text{ smooth.}$$

Claim $\tilde{L} \simeq \mathbb{A}^1 \Rightarrow \tilde{L} \cap \pi^{-1}(0)$ is a single point in $\pi^{-1}(0)^{\tilde{\theta}}$


- The lift $\tilde{\theta}$ sends $\pi^{-1}(0)$ to $\pi^{-1}(0)$ b/c ($\theta(0)=0$),
by permuting the components $C_i \simeq \mathbb{P}^1$. (swapping or preserve)

Lemma ^{Exercise.}: An involution on \mathbb{P}^1 either acts trivially or has exactly two fixed points.

(comes from Möbius transformation)


Example (continued):

Type A: $\theta: x \leftrightarrow y$
n odd



$\pi^{-1}(0)^{\tilde{\theta}} =$ two discrete points
action of $\tilde{\theta}$


n even



$\pi^{-1}(0)^{\tilde{\theta}} =$ a discrete point

§ 3-2 Nakajima quiver variety

Example: Nakajima quiver variety type A.



Attach to each vertex a 1-dim'l vector space V_i

$\dim V_i = m_i$ ($= 1$ in type A)
 $m_0 = 1$ coefficient in $\mathfrak{g} = \sum_{i=1}^n m_i \alpha_i$

Consider the representation space

$$M(\mathcal{V}) = \mathbb{T}^* \left(\bigoplus_{i=0}^n \text{Hom}(V_i, V_{i+1}) \right) = \bigoplus_{i=0}^n \text{Hom}(V_i, V_{i+1}) \oplus \bigoplus_{i=0}^n \text{Hom}(V_{i+1}, V_i)$$

symplectic vector space

$$G = \prod_{i=0}^n GL(V_i) \curvearrowright M(\mathcal{V}, \omega)$$

moment map $\mu = (\mu_i)_{i=0}^n : \mu_i = a_i b_{i-1} - b_i a_i$

We have $\mu^{-1}(0) // G = \text{Spec } \mathbb{C}[\underbrace{\mu^{-1}(0)}_{\text{is also graded}}] // G \simeq \text{Spec } \mathbb{C}[x, y, z] / (xy - z^{n+1})$
 Type A_n Kleinian singularity.

with $x := a_n a_{n-1} \cdots a_1 a_0$, $y := b_0 b_1 \cdots b_{n-1} b_n$, $z := a_0 b_0$ #

(product of determinant maps)

Take $\chi: G \rightarrow \mathbb{C}^*$, a generic character.

consider $\mu^{-1}(0)^S \subset \mu^{-1}(0)$ (semi) stable locus. $\sim G^2 \mu^{-1}(0)^S$ freely

Then $\tilde{X} \cong \mu^{-1}(0)^S // G \longrightarrow \mu^{-1}(0) // G \cong X$

is the minimal resolution of Kleinian singularities [Kroheimer 89, Nakajima 94]

Example (continued)

(1) $M(V) \rightarrow M(V)$

(2) $(a_i) = b_{n-2}$ (1) $(b_i) = a_{n-2}$, anti-symplectic involution

- then
- (1) (1) preserves $\mu^{-1}(0)$
 - (2) (1) preserves $\mu^{-1}(0)^S$
 - (3) (1) normalizes G

routine check

(4) (4) $(x) = \gamma$, (4) $(\gamma) = x$, (4) $(z) = z \sim$ (4) descends to Θ on X .

Thm (Nakajima 96) $C_i = \{ (a, b) \in \mu^{-1}(0)^S // G \mid a_i = b_{i-1} = 0 \}$

so $\Theta(C_i) = C_{n+1-i}$ (swapping the components C_i, C_{n+1-i})
 corresponds to folding of Dynkin diagram A_n ,

Rmk: types D, E can be done similarly; but main difficulty is that:
 $\dim V_i = M_i$, not just 1-dim! making it hard to find invariant functions x, y, z.
 & no triple intersection in types D, E

§ 3.3 multiplicities

X^θ irreducible components L_1, \dots, L_m
reduced

$\pi^{-1}(X^\theta)$ irreducible components $\underbrace{\tilde{L}_1, \dots, \tilde{L}_m}_{|A|}, \underbrace{C_1, \dots, C_n}_{|P|}$

$$\pi^{-1}(X^\theta) = \sum_{j=1}^m \tilde{L}_j + \sum_{i=1}^n \boxed{a_i} C_i \text{ as a divisor.}$$

↑
want to determine a_i

Define $b_i := \#\{ \tilde{L}_j \mid \tilde{L}_j \cap C_i \neq \emptyset \}$

Prop 4 (multiplicity) If $X^\theta \subset X$ is a principal divisor,


then
$$\begin{pmatrix} a_1 \\ \vdots \\ a_n \end{pmatrix} = \boxed{e^{-1}} \begin{pmatrix} b_1 \\ \vdots \\ b_n \end{pmatrix}.$$

inverse of
Cartan matrix

Rmk: $X^\theta \subset X$ is principal modulo two cases \rightarrow have other method to determine multiplicity

Example (continued)


Type A_n . $\theta: X \hookrightarrow Y$ (case 1)



$b_1 = 0, b_2 = 2, b_3 = 0$

$$e \begin{pmatrix} a_1 \\ a_2 \\ a_3 \end{pmatrix} = \begin{pmatrix} b_1 \\ b_2 \\ b_3 \end{pmatrix} \rightsquigarrow \begin{pmatrix} 2 & -1 & 0 \\ -1 & 2 & -1 \\ 0 & -1 & 2 \end{pmatrix} \begin{pmatrix} a_1 \\ a_2 \\ a_3 \end{pmatrix} = \begin{pmatrix} 0 \\ 2 \\ 0 \end{pmatrix}$$

$\Rightarrow a_1 = 1, a_2 = 2, a_3 = 1$



Not reduced unless in type A_1 .